

# MOMENT OPERATORS OF THE CARTESIAN MARGINS OF THE PHASE SPACE OBSERVABLES

J. KIUKAS, P. LAHTI, AND K. YLINEN

**ABSTRACT.** The theory of operator integrals is used to determine the moment operators of the Cartesian margins of the phase space observables generated by the mixtures of the number states. The moments of the  $x$ -margin are polynomials of the position operator and those of the  $y$ -margin are polynomials of the momentum operator.

## 1. INTRODUCTION

According to the theory developed in [7], each complex valued measurable function  $f$  and operator measure  $E$  determine a unique, possibly unbounded, linear operator  $L(f, E)$ , the operator integral of  $f$  with respect to  $E$ . In the case of real valued functions and phase space operator measures, a natural application of this theory is quantization.

In general, quantization means any procedure which associates a quantum mechanical observable to a given classical dynamical variable, the latter being represented by a real valued measurable function on the phase space  $\mathbb{R}^{2n}$  of the classical system. Phase space quantization schemes are often realized by associating to a given classical variable  $f$  the operator  $\int f(q, p) \Delta(q, p) dq dp$ , where  $\Delta$  is some operator valued phase space function and the integral is interpreted e.g. in the weak or strong sense on a suitable domain (cf. e.g. [6], [9], and [13]).

Consider the quantization determined by the operator density  $(q, p) \mapsto \Delta(q, p) := (2\pi)^{-1} W(-q, p) T W(-q, p)^*$ , where the  $W(-q, p)$  are the Weyl operators acting in a separable Hilbert space and  $T$  is a state, i.e. a positive operator of trace one. Now the map  $\mathcal{B}(\mathbb{R}^{2n}) \ni B \mapsto E(B) := \int_B \Delta \in L(\mathcal{H})$  is a phase space observable, so that any classical variable  $f$  can be integrated with respect to it. On its natural domain, the operator integral  $L(f, E)$  coincides with the (weak) quantization integral  $\int f \Delta$ . In this way,  $L(f, E)$  can be interpreted as a quantization of the classical variable  $f$ . It can be noted that this approach differs from the Weyl quantization, which is obtained by replacing  $T$  in the above density  $\Delta(q, p)$  by (a constant times) the parity operator in  $L^2(\mathbb{R})$  (cf. e.g. [3, Sect. IV.1], [6, p. 199], and [9, pp. 140-141]).

In this paper, we consider the phase space observables on  $\mathbb{R}^2$  associated with certain pure states and their convex combinations, with the state vectors being taken from a fixed countable orthonormal basis of the separable Hilbert

space. The moment operators of the Cartesian margins of these operator measures will be determined using the theory of operator integrals. The results sharpen and extend some of those obtained previously in [8].

sectionOperator integrals and phase space observables

**1.1. The operator integral.** In the following, we review the basic results of [7] on the theory of operator integrals and prove a proposition concerning integration with respect to sequences of positive operator measures.

Let  $\mathcal{H}$  be a Hilbert space, with inner product  $\langle \cdot | \cdot \rangle$ , and  $L(\mathcal{H})$  the set of bounded operators on  $\mathcal{H}$ . Let  $\Omega$  be a nonempty set,  $\mathcal{A}$  a  $\sigma$ -algebra of subsets of  $\Omega$ , and  $E : \mathcal{A} \rightarrow L(\mathcal{H})$  a positive operator measure (a positive operator valued set function  $\sigma$ -additive with respect to the strong, or, equivalently, the weak operator topology). For all  $\varphi, \psi \in \mathcal{H}$ , the map  $\mathcal{A} \ni X \mapsto \langle \psi | E(X) \varphi \rangle \in \mathbb{C}$  is a complex measure, and it is denoted by  $E_{\psi, \varphi}$ . Let  $f : \Omega \rightarrow \mathbb{C}$  be an  $\mathcal{A}$ -measurable function and let  $D(f, E)$  be the set of those vectors  $\varphi \in \mathcal{H}$  for which  $f$  is integrable with respect to the complex measure  $E_{\psi, \varphi}$  for all  $\psi \in \mathcal{H}$ . The set  $D(f, E)$  is a vector subspace of  $\mathcal{H}$ , and there is a unique linear operator  $L(f, E)$  on the domain  $D(f, E)$  such that

$$\langle \psi | L(f, E) \varphi \rangle = \int f dE_{\psi, \varphi}$$

for all  $\varphi \in D(f, E)$  and  $\psi \in \mathcal{H}$  (cf. [7]). We call  $L(f, E)$  the (*operator*) *integral* of  $f$  with respect to  $E$ .

Let  $\tilde{D}(f, E)$  be the set of those vectors  $\varphi \in \mathcal{H}$  for which  $|f|^2$  is integrable with respect to the positive measure  $E_{\varphi, \varphi}$ . We have the following results, proved in [7]:

- Theorem 1.** (a)  $\tilde{D}(f, E)$  is a vector subspace of  $D(f, E)$ .  
 (b) If  $E(X)$  is a projection for all  $X \in \mathcal{A}$ , then  $\tilde{D}(f, E) = D(f, E)$ .

It is well known that, in case (b), the domain is dense.

**Theorem 2.** If  $f$  is real valued, then  $L(f, E)$  is a symmetric operator, that is,  $\langle \psi | L(f, E) \varphi \rangle = \langle L(f, E) \psi | \varphi \rangle$  for all  $\psi, \varphi \in D(f, E)$ .

The following general lemma will be used in the proof of Proposition 1.

**Lemma 1.** Let  $\mu_n : \mathcal{A} \rightarrow \mathbb{C}$  be a complex measure for each  $n \in \mathbb{N}$  such that the series  $\sum_{n \in \mathbb{N}} \mu_n$  converges absolutely in the total variation norm. Let  $\mu$  and  $\nu$  denote the measures  $\sum_{n \in \mathbb{N}} \mu_n$  and  $\sum_{n \in \mathbb{N}} |\mu_n|$ , respectively. Here  $|\cdot|$  stands for the total variation.

- (a)  $f$  is  $\nu$ -integrable, if and only if  $\sum_{n=1}^{\infty} \int |f| d|\mu_n| < \infty$ .  
 (b) If  $f$  is  $\nu$ -integrable, then  $f$  is integrable with respect to  $\mu$  and all the measures  $\mu_n$ , and

$$\int f d\mu = \sum_{n=1}^{\infty} \int f d\mu_n.$$

**Proof.** (a) Assume that  $f$  is  $\nu$ -integrable (i.e.  $|f|$  is such). Since  $|\mu_n(B)| \leq \nu(B)$  for all  $B \in \mathcal{A}$  and  $n \in \mathbb{N}$ ,  $|f|$  is  $\mu_n$ -integrable for each  $n \in \mathbb{N}$ . Now

$$\sum_{n=1}^k \int |f| d|\mu_n| = \int |f| d\left(\sum_{n=1}^k |\mu_n|\right) \leq \int |f| d\nu$$

for each  $k \in \mathbb{N}$ , so that  $\sum_{n=1}^{\infty} \int |f| d|\mu_n| < \infty$ . The converse follows from Lemma A.5. of [7].

(b) Clearly  $\mu$  and all the  $\mu_n$  are  $\nu$ -continuous. Let  $g$  and  $g_n$  be their Radon-Nikodým derivatives with respect to  $\nu$ , respectively. Since

$$\left(\sum_{n=1}^k \mu_n - \mu\right)(B) = \int_B \left(\sum_{n=1}^k g_n - g\right) d\nu$$

for all  $B \in \mathcal{A}$  and  $k \in \mathbb{N}$ , we have

$$\lim_{k \rightarrow \infty} \int \left| \sum_{n=1}^k g_n - g \right| d\nu = \lim_{k \rightarrow \infty} \left\| \sum_{n=1}^k \mu_n - \mu \right\| = 0,$$

where  $\|\cdot\|$  denotes the total variation norm. This means that the series  $\sum_n g_n$  converges in  $L^1(\nu)$  to the function  $g$ . Thus some subsequence  $\left(\sum_{n=1}^{k_m} g_n\right)$  converges to  $g$   $\nu$ -almost everywhere. In addition, the monotone convergence theorem gives

$$\nu(B) = \sum_{n=1}^{\infty} \int_B |g_n| d\nu = \int_B \left(\sum_n |g_n|\right) d\nu$$

for all  $B \in \mathcal{A}$ , so that

$$(1) \quad \left| f(x) \sum_{n=1}^{k_m} g_n(x) \right| \leq |f(x)| \sum_n |g_n| = |f(x)|$$

for  $\nu$ -almost all  $x \in \Omega$ .

Assume now that  $f$  is  $\nu$ -integrable. Then by (a),  $f$  is  $\mu_n$ -integrable for each  $n \in \mathbb{N}$ , and the series  $\sum_{n=1}^{\infty} \int f d\mu_n$  converges absolutely. Because of (1), the dominated convergence theorem implies that  $fg$  is  $\nu$ -integrable, i.e.  $f$  is  $\mu$ -integrable, and

$$\int f d\mu = \int fg d\nu = \lim_{m \rightarrow \infty} \sum_{n=1}^{k_m} \int f g_n d\nu = \sum_{n=1}^{\infty} \int f d\mu_n.$$

□

**Proposition 1.** Let  $E^n : \mathcal{A} \rightarrow L(\mathcal{H})$  be a positive operator measure for each  $n \in \mathbb{N}$  such that the range of  $E^n$  is bounded in norm by  $M_n > 0$ , with  $\sum_n M_n < \infty$ . Then the norm limit  $\sum_n E^n(B)$ ,  $B \in \mathcal{A}$ , defines a positive operator measure  $E$ , for which

$$L(f, E)|_{\tilde{D}(f, E)} \subset \sum_n L(f, E^n)|_{\tilde{D}(f, E^n)},$$

where the series is understood to converge in the weak sense to an operator, whose domain consists of those vectors  $\varphi \in \bigcap_{n \in \mathbb{N}} \tilde{D}(f, E^n)$  for which the vector series  $\sum_n L(f, E^n)\varphi$  converges weakly.

**Proof.** Let  $\psi, \varphi \in \mathcal{H}$ . Since  $\|E_{\psi, \varphi}^n\| \leq 4 \sup_{B \in \mathcal{A}} |E_{\psi, \varphi}^n(B)| \leq 4\|\psi\|\|\varphi\|M_n$ , the series  $\sum_n E_{\psi, \varphi}^n$  converges absolutely in the total variation norm. The above inequality also implies that for every  $B \in \mathcal{A}$ , the sesquilinear functional  $(\varphi, \psi) \mapsto \sum_n E_{\psi, \varphi}^n(B)$  is bounded, so that there is a (clearly positive) operator  $E(B) \in L(\mathcal{H})$ , for which  $\langle \psi | E(B) \varphi \rangle = \sum_n E_{\psi, \varphi}^n(B)$ . Thus the map  $B \mapsto E(B)$  is a positive operator measure, for which  $E(B) = \sum_n E^n(B)$  in the operator norm and  $E_{\psi, \varphi} = \sum_n E_{\psi, \varphi}^n$  in the total variation norm.

By applying Lemma 1 (b) to the finite positive measures  $E_{\varphi, \varphi}^n$  and the function  $|f|^2$ , we see that if  $\varphi \in \tilde{D}(f, E)$ , then  $\varphi \in \tilde{D}(f, E^n)$  for all  $n \in \mathbb{N}$ , and  $\sum_n \int |f|^2 dE_{\varphi, \varphi}^n = \int |f|^2 dE_{\varphi, \varphi} < \infty$ . Now let  $F^n$  denote the positive operator measure  $M_n^{-1}E^n$ . We have  $\sum_n M_n \sqrt{\int |f|^2 dF_{\psi, \varphi}^n} \leq \sum_n M_n (1 + \int |f|^2 dF_{\psi, \varphi}^n) < \infty$ . Using the inequality

$$\int |f| d|F_{\psi, \varphi}^n| \leq \|\psi\| \sqrt{\int |f|^2 dF_{\varphi, \varphi}^n}$$

from [7], we get  $\sum_n \int |f| d|E_{\psi, \varphi}^n| = \sum_n M_n \int |f| d|F_{\psi, \varphi}^n| < \infty$ , from which it follows by lemma 1 (a) and (b) that

$$\langle \psi | L(f, E) \varphi \rangle = \sum_n \langle \psi | L(f, E^n) \varphi \rangle.$$

□

**1.2. Phase space observables.** We assume that the Hilbert space  $\mathcal{H}$  is separable. For any  $\varphi, \psi \in \mathcal{H}$ , let  $|\varphi\rangle\langle\psi|$  denote the operator  $\xi \mapsto \langle\psi|\xi\rangle\varphi$ . Let  $\{|n\rangle \mid n \geq 0\}$  be a fixed orthonormal basis of  $\mathcal{H}$ . We call it the number basis. Let  $A_+ = \sum_{n \geq 0} \sqrt{n+1}|n+1\rangle\langle n|$  and  $A_- = \sum_{n \geq 0} \sqrt{n+1}|n\rangle\langle n+1|$  be the raising and lowering operators associated with this basis. They are closed operators with the domain

$$\mathcal{D}(A_+) = \mathcal{D}(A_-) = \left\{ \varphi \in \mathcal{H} \mid \sum_k k |\langle \varphi | k \rangle|^2 < \infty \right\},$$

and they satisfy the relation  $A_+ = A_-^*$ . The symmetric operators  $\frac{1}{\sqrt{2}}(A_+ + A_-)$  and  $\frac{1}{\sqrt{2}}i(A_+ - A_-)$  are essentially selfadjoint on  $\mathcal{D}(A_+)$  and their closures  $Q$  and  $P$  are unitarily equivalent to the canonical position and momentum operators, respectively, acting in  $L^2(\mathbb{R})$ . The operators  $A_+$  and  $A_-$  can be

expressed in terms of  $Q$  and  $P$  according to

$$\begin{aligned} A_+ &= \frac{1}{\sqrt{2}}(Q - iP), \\ A_- &= \frac{1}{\sqrt{2}}(Q + iP) \end{aligned} \quad (2)$$

(cf. [2, p. 283] and [12, p. 73]). The number operator  $N := \sum_{n \geq 0} n|n\rangle\langle n| = A_+A_-$ , with the domain

$$D(N) = \left\{ \varphi \in \mathcal{H} \mid \sum_k k^2 |\langle \varphi | k \rangle|^2 < \infty \right\},$$

is selfadjoint and satisfies

$$N + \frac{1}{2}I = \frac{1}{2}(Q^2 + P^2). \quad (3)$$

(The last equality is a simple consequence of (2) and the fact that  $D(N) = D(A_+A_-) = D(A_-A_+)$ .)

The operators  $Q$  and  $P$  generate strongly continuous one parameter unitary groups  $p \mapsto V(p) := e^{ipQ}$  and  $q \mapsto U(q) := e^{iqP}$  which satisfy the Weyl relation  $U(q)V(p) = e^{iqp}V(p)U(q)$  for all  $q, p \in \mathbb{R}$ . The Weyl operators  $W(q, p)$  are defined by  $W(q, p) = e^{-\frac{1}{2}iqp}U(q)V(p)$  for  $q, p \in \mathbb{R}$ . The map  $(q, p) \mapsto W(-q, p)$  is a projective representation of  $\mathbb{R}^2$ .

Let  $T$  be a state operator. Then

$$I = \frac{1}{2\pi} \int_{\mathbb{R}^2} W(-q, p)TW(-q, p)^* dq dp,$$

and the map  $E^T : \mathcal{B}(\mathbb{R}^2) \rightarrow L(\mathcal{H})$  defined by

$$E^T(B) = \frac{1}{2\pi} \int_B W(-q, p)TW(-q, p)^* dq dp$$

is a normalized positive operator measure, a phase space observable. Here  $\mathcal{B}(\mathbb{R}^2)$  denotes the  $\sigma$ -algebra of Borel subsets of  $\mathbb{R}^2$  and the integrals are understood in the weak sense. The construction of the operator measures  $E^T$  can be found for instance in [5] or [14]. The operator measure  $E^T$  is covariant with respect to the projective representation  $(q, p) \mapsto W(-q, p)$  in the sense that  $E^T(B + (q_0, p_0)) = W(-q_0, p_0)E^T(B)W(-q_0, p_0)^*$  for all  $B \in \mathcal{B}(\mathbb{R}^2)$  and  $(q, p) \in \mathbb{R}^2$ .

The following characterization is obtained in [4] and [15]: every normalized positive operator measure  $E : \mathcal{B}(\mathbb{R}^2) \rightarrow L(\mathcal{H})$ , which is covariant with respect to the representation  $(q, p) \mapsto W(-q, p)$ , is of the form  $E^T$  for some state  $T$ .

We let  $E^{|s\rangle}$  denote the phase space observable associated with the number state  $|s\rangle\langle s|$ . Consider the mixed states

$$T = \sum_{n=0}^{\infty} w_n |n\rangle\langle n|, \quad (4)$$

where  $w_n \geq 0$  and  $\sum w_n = 1$ . These states are the ones for which the observable  $E^T$  is covariant also with respect to the phase shifts in the sense that

$$e^{i\theta N} E^T([0, \infty) \times X) e^{-i\theta N} = E^T([0, \infty) \times (X + \theta))$$

for all  $\theta \in [0, 2\pi)$  and  $X \in \mathcal{B}([0, 2\pi))$ , where  $\mathbb{R}^2 = [0, \infty) \times [0, 2\pi)$  and the sum  $X + \theta$  is understood modulo  $2\pi$ . (cf. [11]).

Since  $w_n \|E^{[n]}(B)\| \leq w_n$  for all  $n \in \mathbb{N}$  and  $B \in \mathcal{B}(\mathbb{R}^2)$ , Proposition 1 can be applied to the positive operator measures  $w_n E^{[n]}$ . That the norm limit  $\sum_n w_n E^{[n]}(B)$  equals  $E^T(B)$ , follows from the identity

$$\langle \varphi | E^T(B) \varphi \rangle = \frac{1}{2\pi} \int_B \left( \sum_n w_n |\langle \varphi | W(-q, p) | n \rangle|^2 \right) dq dp = \sum_n w_n \langle \varphi | E^{[n]}(B) \varphi \rangle$$

where  $\varphi \in \mathcal{H}$  is arbitrary and the monotone convergence theorem has been used.

## 2. MOMENT OPERATORS OF THE CARTESIAN MARGINS OF THE PHASE SPACE OBSERVABLES ASSOCIATED WITH THE NUMBER STATES

Let  $x$  and  $y$  denote the functions  $(q, p) \mapsto q$  and  $(q, p) \mapsto p$ , respectively. In [7], the moment operators  $L((x \pm iy)^k, E^{[n]})$  and  $L((x^2 + y^2)^k, E^{[n]})$  were determined. In [8], these results were used to obtain the operator relations

$$\begin{aligned} L(x, E^{[n]}) &\subset Q, \\ L(y, E^{[n]}) &\subset P, \\ L(x^2, E^{[n]}) &\subset (n + \frac{1}{2})I + Q^2, \\ (5) \quad L(y^2, E^{[n]}) &\subset (n + \frac{1}{2})I + P^2. \end{aligned}$$

In this section, we determine directly the moment operators  $L(x^k, E^T)$  and  $L(y^k, E^T)$ , where the state  $T$  is taken to be of the form  $\sum_n w_n |n\rangle\langle n|$ . The results show, among other things, that the inclusions (5) are in fact equalities.

**2.1. The operators  $L(x^k, E^{[n]})$  and  $L(y^k, E^{[n]})$ .** Let  $U : L^2(\mathbb{R}) \rightarrow \mathcal{H}$  be the unitary operator which maps the Hermite function basis  $\{h_n\}_{n \geq 0}$  of  $L^2(\mathbb{R})$  onto the number basis of  $\mathcal{H}$  according to the rule  $U h_n = |n\rangle$ . The position and momentum operators in  $L^2(\mathbb{R})$  correspond to the operators  $Q$  and  $P$  via the unitary transformation  $U$ , so that the operators  $W_0(q, p)$ , defined by  $W_0(q, p) = U^{-1} W(q, p) U$ , act in  $L^2(\mathbb{R})$  according to the formula

$$(W_0(q, p)f)(t) = e^{i\frac{1}{2}qp} e^{ipt} f(t + q).$$

We need the following well-known result (see e.g. [14, pp. 47 and 49]).

**Lemma 2.** *Let  $F : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  denote the Fourier-Plancherel operator. Let  $u \in L^2(\mathbb{R})$  be a unit vector and  $f \in L^2(\mathbb{R})$ . Then  $\overline{u}(\cdot - q)f \in L^1(\mathbb{R}) \cap$*

$L^2(\mathbb{R})$  for almost all  $q \in \mathbb{R}$  and  $\overline{Fu}(\cdot - p)Ff \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$  for almost all  $p \in \mathbb{R}$ . In addition,

$$\frac{1}{\sqrt{2\pi}} \langle W_0(-q, p)u|f \rangle = e^{i\frac{1}{2}qp} F(\overline{u}(\cdot - q)f)(p)$$

for almost all  $q \in \mathbb{R}$  and all  $p \in \mathbb{R}$ , and

$$\frac{1}{\sqrt{2\pi}} \langle W_0(-q, p)u|f \rangle = e^{-i\frac{1}{2}qp} F^{-1}(\overline{Fu}(\cdot - p)Ff)(q)$$

for almost all  $p \in \mathbb{R}$  and all  $q \in \mathbb{R}$ .

**Proof.** Since  $u, f \in L^2(\mathbb{R})$ , the function  $\overline{u}(\cdot - q)f$  is integrable. Because  $\|f\|^2 = \int |\overline{u}(t - q)f(t)|^2 dt dq$  by Fubini's theorem,  $\overline{u}(\cdot - q)f \in L^2(\mathbb{R})$  for almost all  $q$ . Similarly,  $\overline{Fu}(\cdot - p)Ff \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$  for almost all  $p \in \mathbb{R}$ . The rest of the proof follows from straightforward calculations.  $\square$

First we determine the square integrability domains corresponding to the functions  $x^k$  and  $y^k$ .

**Lemma 3.**  $\tilde{D}(x^k, E^{[n]}) = D(Q^k)$  and  $\tilde{D}(y^k, E^{[n]}) = D(P^k)$  for all  $k \in \mathbb{N}$ .

**Proof.** Let  $k \in \mathbb{N}$  be fixed. Let  $\varphi \in \mathcal{H}$  and  $f = U^{-1}\varphi \in L^2(\mathbb{R})$ . If  $\varphi \in \tilde{D}(x^k, E^{[n]})$ , the function  $(q, p) \mapsto q^{2k} |\langle f|W_0(-q, p)h_n \rangle|^2 = q^{2k} |\langle \varphi|W(-q, p)|n \rangle|^2$  is integrable over  $\mathbb{R}^2$ , and

$$\begin{aligned} \int_{\mathbb{R}^2} q^{2k} dE_{\varphi, \varphi}^{[n]}(q, p) &= \frac{1}{2\pi} \int q^{2k} \left( \int |\langle \varphi|W(-q, p)|n \rangle|^2 dp \right) dq \\ &= \int q^{2k} \left( \int |F(\overline{h_n}(\cdot - q)f)(p)|^2 dp \right) dq \\ &= \int q^{2k} \left( \int |h_n(t - q)|^2 |f(t)|^2 dt \right) dq \\ &= \int \left( \int q^{2k} |h_n(t - q)|^2 |f(t)|^2 dq \right) dt \\ &= \int \int (t - q)^{2k} |h_n(q)|^2 |f(t)|^2 dq dt \\ &= \int \int (t - q)^{2k} |h_n(q)|^2 |f(t)|^2 dt dq, \end{aligned}$$

where lemma 2, the unitarity of the Fourier-Plancherel operator, and Fubini's theorem have been used. The existence of the last integral implies that  $t \mapsto (t - q)^{2k} |f(t)|^2$  is integrable over  $\mathbb{R}$  for almost all  $q \in \mathbb{R}$ . Thus also  $t \mapsto t^{2k} |f(t)|^2$  must be integrable. (In fact, take one  $q \in \mathbb{R}$  for which  $t \mapsto (t - q)^{2k} |f(t)|^2$  is integrable and use the fact that there exist positive constants  $A, B, M$ , such that  $At^{2k} \leq (t - q)^{2k} \leq Bt^{2k}$  for  $|t| \geq M$ .) This means that  $f$  belongs to the domain of the  $k$ -th power of the position operator in  $L^2(\mathbb{R})$  and hence  $\varphi = Uf \in D(Q^k)$ . Conversely, if  $\varphi = Uf \in D(Q^k)$ , the functions  $t \mapsto |t|^l |f(t)|^2$  and  $q \mapsto |q|^l |h_n(q)|^2$  are integrable over  $\mathbb{R}$  for

all  $l \leq 2k$  and hence  $(t, q) \mapsto (t - q)^{2k} |h_n(q)|^2 |f(t)|^2$  is integrable over  $\mathbb{R}^2$ . The preceding calculation now shows that  $\varphi \in \tilde{D}(x^k, E^{[n]})$ . The equality  $\tilde{D}(x^k, E^{[n]}) = D(Q^k)$  is thus proved.

The result  $\tilde{D}(y^k, E^{[n]}) = D(P^k)$  is obtained in an analogous manner by using the fact that the position and momentum operators in  $L^2(\mathbb{R})$  are unitarily equivalent via the Fourier-Plancherel operator  $F$ .  $\square$

Now we can determine the operators  $L(x^k, E^{[n]})$  and  $L(y^k, E^{[n]})$ .

**Theorem 3.**  $L(x^k, E^{[n]}) = p_k^{[n]}(Q)$  and  $L(y^k, E^{[n]}) = p_k^{[n]}(P)$ , where  $p_k^{[n]} : \mathbb{R} \rightarrow \mathbb{R}$  is the polynomial

$$p_k^{[n]}(t) = \langle n | (t - Q)^k | n \rangle = \sum_{l=0}^k \left( \binom{k}{l} (-1)^{k-l} \langle n | Q^{k-l} | n \rangle \right) t^l.$$

**Proof.** Since  $p_k^{[n]}$  is a polynomial of order  $k$  and  $Q$  is unitarily equivalent to the position operator in  $L^2(\mathbb{R})$ , the natural domain of the operator  $p_k^{[n]}(Q)$  (which is the set  $D(Q^k) \cap D(Q^{k-1}) \cap \dots \cap D(Q)$ ) is equal to that of  $Q^k$ . Because  $Q$  and  $P$  are unitarily equivalent, also  $D(p_k^{[n]}(P)) = D(P^k)$ . Thus by the preceding lemma, we have  $D(p_k^{[n]}(Q)) = D(Q^k) = \tilde{D}(x^k, E^{[n]})$  and  $D(p_k^{[n]}(P)) = D(P^k) = \tilde{D}(y^k, E^{[n]})$ .

Let  $\varphi, \psi \in \tilde{D}(x^k, E^{[n]}) \subset D(x^k, E^{[n]})$ . Let  $f = U^{-1}\varphi$ ,  $g = U^{-1}\psi$ . Since the function

$$(q, p) \mapsto q^k \langle \psi | W(-q, p) | n \rangle \overline{\langle \varphi | W(-q, p) | n \rangle}$$

is integrable over  $\mathbb{R}^2$ , we get

$$\begin{aligned} \langle \psi | L(x^k, E^{[n]}) \varphi \rangle &= \int_{\mathbb{R}^2} q^k dE_{\psi, \varphi}^{[n]}(q, p) \\ &= \frac{1}{2\pi} \int q^k \left( \int \langle \psi | W(-q, p) | n \rangle \overline{\langle \varphi | W(-q, p) | n \rangle} dp \right) dq \\ &= \int q^k \left( \int \overline{F(\overline{h_n(\cdot - q)g}(p))} F(\overline{h_n(\cdot - q)f}(p)) dp \right) dq \\ &= \int q^k \left( \int \overline{h_n(t - q)g(t)} h_n(t - q)f(t) dt \right) dq \\ &= \int \left( \int q^k |h_n(t - q)|^2 dq \right) \overline{g(t)} f(t) dt \\ &= \int \left( \int (t - q)^k |h_n(q)|^2 dq \right) \overline{g(t)} f(t) dt, \\ &= \int \langle n | (t - Q)^k | n \rangle \overline{g(t)} f(t) dt \\ &= \langle \psi | p_k^{[n]}(Q) \varphi \rangle. \end{aligned}$$



The fifth equality follows from Fubini's theorem, since  $(q, t) \mapsto q^k |h_n(t - q)|^2 \overline{g(t)} f(t)$  is integrable (because of the inequality

$$\left| q^k |h_n(t - q)|^2 \overline{g(t)} f(t) \right| \leq \frac{1}{2} (1 + q^{2k}) (|f(t)|^2 + |g(t)|^2) |h_n(t - q)|^2$$

and the proof of lemma 3). The unitarity of  $F$  is also used. Since  $\psi$  was taken arbitrarily from the dense set  $D(Q^k) = \tilde{D}(f, E^{[n]})$ , we have  $p_k^{[n]}(Q) \subset L(x^k, E^{[n]})$ .

The equality  $p_k^{[n]}(Q) = L(x^k, E^{[n]})$  follows from the fact that being self-adjoint, the operator  $p_k^{[n]}(Q)$  cannot have a proper symmetric extension.

The statement  $p_k^{[n]}(P) = L(y^k, E^{[n]})$  is obtained in the same manner, since  $p_k^{[n]}$  can also be written in the form  $p_k^{[n]}(t) = \langle n | (t - P)^k | n \rangle$  and  $p_k^{[n]}(P)$  is selfadjoint.  $\square$

**Remark.** Since  $\langle n | Q^m | n \rangle = 0$  for odd  $m$ , and  $\langle n | Q^m | n \rangle > 0$  for even  $m$ , only the terms with even  $k - l$  are present in the sum defining the polynomial  $p_k^{[n]}$ , and the coefficients of the corresponding  $x^l$  are all strictly positive. In particular,  $L(x^k, E^{[n]}) \neq Q^k$  and  $L(y^k, E^{[n]}) \neq P^k$  for  $k > 1$  and  $n \geq 0$ , reflecting the difference from the Weyl quantization ([6, p. 229]), as well as the fact that the Cartesian margins of  $E^{[n]}$  are not the spectral measures of  $Q$  and  $P$ .

Using Theorem 3, all the operators  $L(x^k, E^{[n]})$  and  $L(y^k, E^{[n]})$  can be written in terms of  $Q$  and  $P$ , respectively. In particular, the first and second moment operators are the following:

$$\begin{aligned} L(x, E^{[n]}) &= Q \\ L(y, E^{[n]}) &= P \\ L(x^2, E^{[n]}) &= \left(n + \frac{1}{2}\right) I + Q^2 \\ L(y^2, E^{[n]}) &= \left(n + \frac{1}{2}\right) I + P^2. \end{aligned} \tag{6}$$

For the special case of  $n = 0$ , these results were already obtained by formal computations in [1, pp. 28-29] (without addressing the question on the domains of the operators). A related result of [9, p. 140], however, seems to lack a constant term.

**2.2. The operators  $L(x^k, E^T)$  and  $L(y^k, E^T)$  with  $T = \sum_n w_n |n\rangle\langle n|$ .** In the next theorem, we need to consider the expressions  $\langle n | Q^{2k} | n \rangle$ . These integrals are well known and can be calculated e.g. as instructed in [10, p. 60]. We need here only the following fact:

**Lemma 4.** *For  $n \geq k$ , the expression  $\langle n | Q^{2k} | n \rangle$  can be written as a polynomial in  $n$  of order  $k$ .*

**Proof.** Expressing  $Q$  in terms of  $A_+$  and  $A_-$ , we get  $\langle n|Q^{2k}|n\rangle = \frac{1}{2^k} \|(A_+ + A_-)^k|n\rangle\|^2$ . Because  $A_-A_+ = N + I$ , we can write

$$(A_+ + A_-)^k|n\rangle = \sum_{m=0}^k a_m|n+k-2m\rangle,$$

where  $a_m|n+k-2m\rangle = A_+^{k-2m}q_m^+(N)|n\rangle$  for  $0 \leq m \leq \frac{k}{2}$  and  $a_m|n+k-2m\rangle = A_-^{2m-k}q_m^-(N)|n\rangle$  for  $\frac{k}{2} \leq m \leq k$ , where  $q_m^\pm$  are some polynomials with  $2\deg(q_m^\pm) \pm (k-2m) \leq k$ . Now  $a_m^2 = (n+1)(n+2)\dots(n+k-2m)q_m^+(n)^2$  for  $0 \leq m \leq \frac{k}{2}$  and  $a_m^2 = n(n-1)\dots(n-(2m-k)+1)q_m^-(n)^2$  for  $\frac{k}{2} \leq m \leq k$ , so that each  $a_m^2$  is a polynomial in  $n$  of order at most  $k$ , with the coefficient of the highest power positive. Since  $a_0^2 = (n+1)(n+2)\dots(n+k)$ , we see that  $\langle n|Q^{2k}|n\rangle = \frac{1}{2^k} \sum_{m=0}^k a_m^2$  is a polynomial in  $n$  of order exactly  $k$ .  $\square$

**Theorem 4.** Let  $T = \sum_n w_n|n\rangle\langle n|$  be a mixture of the number states and  $k \in \mathbb{N}$ . Let  $p_k^{[n]}$  denote the polynomials defined in Theorem 3, and define

$$s_{kl} = \binom{k}{l} \sum_{n=0}^{\infty} w_n \langle n|Q^{k-l}|n\rangle \ (\leq \infty)$$

for  $0 \leq l \leq k$ .

(a)  $\tilde{D}(x^k, E^T) \neq \{0\}$  if and only if

$$(7) \quad \sum_n n^k w_n < \infty,$$

in which case  $s_{kl} < \infty$  for all  $0 \leq l \leq k$ ,  $D(x^k, E^T) = \tilde{D}(x^k, E^T) = D(Q^k)$ , and

$$L(x^k, E^T) = \sum_{l=0}^k s_{kl} Q^l.$$

In particular, the operator  $L(x^k, E^T)$  is then selfadjoint.

(b) The statement (a) holds true when "x" and "Q" are replaced by "y" and "P".

**Proof.** Let  $k \in \mathbb{N}$  be fixed. According to Proposition 1, Lemma 3 and Theorem 3 we have,

$$(8) \quad L(x^k, E^T)|_{\tilde{D}(x^k, E^T)} \subset \sum_n w_n L(x^k, E^{[n]}) = \sum_n w_n p_k^{[n]}(Q)$$

(where the series of operators are understood in the same sense as in Proposition 1). Let  $\varphi \in D(Q^k) \supset D(\sum_n w_n p_k^{[n]}(Q))$  and  $\varphi \neq 0$ . Let  $A_{\varphi, \varphi}^{[n]}$  be the density function of the positive measure  $E_{\varphi, \varphi}^{[n]}$ . By the definition of the square integrability domain,  $\varphi \in \tilde{D}(x^k, E^T)$  if and only if the function  $x^{2k} \sum_n w_n A_{\varphi, \varphi}^{[n]}$  is integrable. By the monotone convergence theorem,

the latter statement is equivalent to

$$(9) \quad \sum_n w_n \int x^{2k} dE_{\varphi, \varphi}^{[n]} < \infty.$$

According to the proof of Lemma 3,

$$\begin{aligned} \int x^{2k} dE_{\varphi, \varphi}^{[n]} &= \int \int (t - q)^{2k} |h_n(q)|^2 |(U^{-1}\varphi)(t)|^2 dt dq \\ &= \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{2k-l} \left( \int t^l |(U^{-1}\varphi)(t)|^2 dt \right) \langle n | Q^{2k-l} | n \rangle. \end{aligned}$$

Since  $\langle n | Q^m | n \rangle = 0$  for odd  $m \in \mathbb{N}$ , only the terms with even  $l$  are present in the above sum. Because  $U^{-1}\varphi \neq 0$ , these terms are all strictly positive. Since  $0 < \langle n | Q^{2l} | n \rangle \leq \int (1 + q^{2k}) |h_n(q)|^2 dq = 1 + \langle n | Q^{2k} | n \rangle$  for all  $l \leq k$  and  $\sum_n w_n = 1$ , the convergence of the series  $\sum_n w_n \langle n | Q^{2k} | n \rangle$  implies the convergence of each series  $\sum_n w_n \langle n | Q^{2l} | n \rangle$  and  $s_{kl}$ ,  $l \leq k$ . Thus, it follows that a nonzero vector in  $D(Q^k)$  belongs to  $\tilde{D}(x^k, E^T)$  if and only if the series

$$(10) \quad \sum_n w_n \langle n | Q^{2k} | n \rangle$$

converges. By the preceding lemma, this is equivalent to (7). Since  $\tilde{D}(x^k, E^T) \subset D(Q^k)$  by (8), we have shown that  $\tilde{D}(x^k, E^T) \neq \{0\}$  if and only if (7) holds, and in that case,  $\tilde{D}(x^k, E^T) = D(Q^k)$  and  $s_{kl} < \infty$  for  $l \leq k$ .

From the definition of the polynomials  $p_k^{[n]}$  we see that if (10) converges, then (using (8)) we get

$$\langle \psi | L(x^k, E^T) \varphi \rangle = \sum_n w_n \langle \psi | p_k^{[n]}(Q) \varphi \rangle = \sum_{l=0}^k s_{kl} \langle \psi | Q^l \varphi \rangle$$

for each  $\varphi \in D(Q^k)$  and  $\psi \in \mathcal{H}$ . Thus,  $\sum_{l=0}^k s_{kl} Q^l \subset L(x^k, E^T)$  (now  $D(\sum_{l=0}^k s_{kl} Q^l) = D(Q^k)$ , because  $s_{kk} = 1$ ). Since the operator  $\sum_{l=0}^k s_{kl} Q^l$  is selfadjoint and  $L(x^k, E^T)$  is symmetric, the statement (a) has been proved.

The result (b) is obtained in the same manner, since

$$L(y^k, E^T)|_{\tilde{D}(y^k, E^T)} \subset \sum_n w_n L(y^k, E^{[n]}) = \sum_n w_n p_k^{[n]}(P),$$

$\langle n | Q^m | n \rangle = \langle n | P^m | n \rangle$  for all  $m, n \geq 0$  and

$$\begin{aligned} \int y^{2k} dE_{\varphi, \varphi}^{[n]} &= \int \int (t - p)^{2k} |F h_n(p)|^2 |(FU^{-1}\varphi)(t)|^2 dt dp \\ &= \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{2k-l} \left( \int t^l |(FU^{-1}\varphi)(t)|^2 dt \right) \langle n | P^{2k-l} | n \rangle \end{aligned}$$

for all  $\varphi \in D(P^k)$ .  $\square$

**Remark.** For each  $k \in \mathbb{N}$ , there are states  $T$  of the form (4) such that (7) does not converge (so that  $\tilde{D}(x^k, E^T) = \{0\}$ ), but all the series  $s_{kl}$  do. In that case,  $\sum_{l=0}^k s_{kl} Q^l$  is still a well-defined selfadjoint operator with the domain  $D(Q^k)$ . An example of such a state for  $k \in \mathbb{N}$  is given by  $t_n = \frac{1}{Sn^{k+1}}$ , where  $S = \sum_n n^{-(k+1)}$ . We do not know whether there are any nonzero vectors in the domain  $D(x^k, E^T)$  then.

### 3. OPERATOR INTEGRALS OF SOME POLYNOMIALS

In this last section, we use Theorem 3 to determine the operator integrals for certain types of polynomials. To that end, let  $h, h_1, h_2$  be real polynomials defined by  $h(t) = \sum_{l=0}^k a_l t^l$ ,  $a_k \neq 0$ , and  $h_i(t) = \sum_{l=0}^{k_i} a_{i,l} t^l$ ,  $a_{i,k_i} \neq 0$ ,  $i = 1, 2$ .

**The operators  $L(h \circ x, E^{[n]})$  and  $L(h \circ y, E^{[n]})$ .** Let  $\psi, \varphi \in \mathcal{H}$ . There exist positive constants  $M, A, B$  such that  $A|t^k| \leq |h(t)| \leq B|t^k|$  for  $|t| \geq M$ , which implies that the function  $h \circ x$  (i.e.  $(q, p) \mapsto h(q)$ ) is  $E_{\psi, \varphi}^{[n]}$ -integrable if and only if  $x^k$  is such, and in that case,  $\int h \circ x dE_{\psi, \varphi}^{[n]} = \sum_{l=0}^k a_l \int x^l dE_{\psi, \varphi}^{[n]}$ . Since  $D(x^k, E^{[n]}) = D(Q^k) \subset D(Q^l) = D(x^l, E^{[n]})$  for  $l \leq k$ , it follows that  $D(h \circ x, E^{[n]}) = D(Q^k)$ , and

$$(11) \quad L(h \circ x, E^{[n]}) = \sum_{l=0}^k a_l p_l^{[n]}(Q).$$

Naturally, a similar result holds for the function  $h \circ y$ .

**The operators  $L(h_1 \circ x + ih_2 \circ y, E^{[n]})$ .** Let  $\psi, \varphi \in \mathcal{H}$ . The function  $h_1 \circ x + ih_2 \circ y$  is  $E_{\psi, \varphi}^{[n]}$ -integrable if and only if both  $h_1 \circ x$  and  $h_2 \circ y$  are such, and in that case  $\int (h_1 \circ x + ih_2 \circ y) dE_{\psi, \varphi}^{[n]} = \int h_1 \circ x dE_{\psi, \varphi}^{[n]} + i \int h_2 \circ y dE_{\psi, \varphi}^{[n]}$ . Thus, we have  $D(h_1 \circ x + ih_2 \circ y, E^{[n]}) = D(Q^{k_1}) \cap D(P^{k_2})$ , and

$$(12) \quad L(h_1 \circ x + ih_2 \circ y, E^{[n]}) = \sum_{l=0}^{k_1} a_{1,l} p_l^{[n]}(Q) + i \sum_{l=0}^{k_2} a_{2,l} p_l^{[n]}(P).$$

**The operators  $L(h_1 \circ x + h_2 \circ y, E^{[n]})$  with  $k_i$  even and  $a_{i,k_i} > 0$ .** Assume that  $k_i$  is even, and  $a_{i,k_i} > 0$ ,  $i = 1, 2$ . Then we can choose positive constants  $M, A_i, B_i$  such that  $h_i(t) \geq 0$  and  $A_i t^{k_i} \leq h_i(t) \leq B_i t^{k_i}$  for  $|t| \geq M$  and  $i = 1, 2$ . This implies that the function  $h_1 \circ x + h_2 \circ y$  is  $E_{\psi, \varphi}^{[n]}$ -integrable for  $\psi, \varphi \in \mathcal{H}$  if and only if both  $x^{k_1}$  and  $y^{k_2}$  are such. We get  $D(h_1 \circ x + h_2 \circ y, E^{[n]}) = D(Q^{k_1}) \cap D(P^{k_2})$ , and

$$(13) \quad L(h_1 \circ x + h_2 \circ y, E^{[n]}) = \sum_{l=0}^{k_1} a_{1,l} p_l^{[n]}(Q) + \sum_{l=0}^{k_2} a_{2,l} p_l^{[n]}(P).$$

**A note on the operators  $L(x \pm iy, E^{[n]})$  and  $L(x^2 + y^2, E^{[n]})$ .** The above observations show that, in particular,  $L(\frac{1}{\sqrt{2}}(x \pm iy), E^{[n]}) = \frac{1}{\sqrt{2}}(Q \pm iP)$ , and  $L(\frac{1}{2}(x^2 + y^2), E^{[n]}) = \frac{1}{2}(Q^2 + P^2) + (n + \frac{1}{2})I$ . These operator integrals have already been determined in [7] (using a different method) to be the following:  $L(\frac{1}{\sqrt{2}}(x \pm iy), E^{[n]}) = A_{\mp}$  and  $L(\frac{1}{2}(x^2 + y^2), E^{[n]}) = N + (n + 1)I$ . The fundamental operator equalities (2) and (3) now show that the results are indeed consistent.

**Acknowledgement.** The authors thank Drs. Daniel Dubin and Mark Hennings for pointing out some details concerning the validity of the operator equality (3).

## REFERENCES

- [1] S. T. Ali, J.-P. Antoine, J.-P. Gazeau; Coherent States, Wavelets and Their Generalizations, Springer-Verlag, New York, 2000.
- [2] M. S. Birman, M. Z. Solomjak; Spectral Theory of Self-Adjoint Operators in Hilbert Space, D. Reidel Publishing Co., Dordrecht, 1987.
- [3] P. Busch, M. Grabowski, P. Lahti; Operational Quantum Physics, 2nd Corrected Printing, Springer-Verlag, Berlin, 1997.
- [4] G. Cassinelli, E. De Vito, A. Toigo; Positive operator valued measures covariant with respect to an irreducible representation, J. Math. Phys. **44** (2003) 4768-4775.
- [5] E. B. Davies; Quantum Theory of Open Systems, Academic Press, London, 1976.
- [6] D. A. Dubin, M. A. Hennings, T. B. Smith; Mathematical Aspects of Weyl Quantization and Phase, World Scientific, Singapore, 2000.
- [7] P. Lahti, M. Maczyński, K. Ylinen; The moment operators of phase space observables and their number margins, Rep. Math. Phys. **41** (1998) 319-331.
- [8] P. Lahti, J.-P. Pellonpää, K. Ylinen; Operator integrals and phase space observables, J. Math. Phys. **40** (1999) 2181-2189.
- [9] N. P. Landsman; Mathematical Topics Between Classical and Quantum Mechanics, Springer-Verlag, New York, 1998.
- [10] E. Merzbacher; Quantum Mechanics, Second Edition, John Wiley & Sons, inc., New York, 1970.
- [11] P. Lahti, J.-P. Pellonpää; Covariant phase observables in quantum mechanics, J. Math. Phys. **40** (1999) 4688-4698.
- [12] C. R. Putnam; Commutation Properties of Hilbert Space Operators and Related Topics, Springer-Verlag, Berlin 1967.
- [13] F. E. Schroeck, Jr.; Quantum Mechanics on Phase Space, Kluwer Academic Publishers, Dordrecht, 1996.
- [14] W. Stulpe; Classical Representations of Quantum Mechanics Related to Statistically Complete Observables, Wissenschaft und Technik Verlag, Berlin, 1997.
- [15] R. Werner; Quantum harmonic analysis on phase space, J. Math. Phys. **25** (1984) 1404-1411.

JUKKA KIUKAS, DEPARTMENT OF PHYSICS, UNIVERSITY OF TURKU, FIN-20014  
TURKU, FINLAND

*E-mail address:* `jukka.kiukas@utu.fi`

PEKKA LAHTI, DEPARTMENT OF PHYSICS, UNIVERSITY OF TURKU, FIN-20014 TURKU,  
FINLAND

*E-mail address:* `pekka.lahti@utu.fi`

KARI YLINEN, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TURKU, FIN-20014  
TURKU, FINLAND

*E-mail address:* `kari.ylinen@utu.fi`